

ParaBiotic Plus™

Serving Size 4 capsules

Servings Per Container 30

Amount Per Serving

Wormwood extract (<i>Artemisia annua</i>)	400 mg
Olive leaf extract (<i>Olea europaea</i>) (17-23% oleuropein)	300 mg
Berberine sulfate (from <i>Berberis aquifolium</i> root bark)	200 mg
Grapefruit seed (4:1 extract)	150 mg
Thyme (4:1)	200 mg
Barberry extract (<i>Berberis vulgaris</i>) (6 % berberine)	100 mg
Uva Ursi (4:1 extract)	100 mg
Black Walnut Hull (<i>Juglans nigra</i> , 10:1 extract)	100 mg
Mexican Oregano Extract (<i>Origanum Vulgare</i>)	200 mg

SUGGESTED DOSE: As a dietary supplement, take 1-4 capsules, three times daily, or as directed by your healthcare professional.

PARABIOTIC PLUS™

COMPREHENSIVE SUPPORT FOR HEALTHY INTESTINAL & IMMUNE SYSTEM FUNCTION

- Highly effective botanicals
- Proven ingredients

ParaBiotic Plus™ is an excellent choice in the nutraceutical management of infection in your patient population. It is safe for use in adolescents over age 12, and adults. This formula is not recommended for use during pregnancy, by nursing mothers, or by children under the age of 12. ParaBiotic Plus™ is a safe, non-prescription alternative for infections of many types. The ingredients in ParaBiotic Plus™ have been proven over decades of clinical use and shown effective in clinical trials. Many nutritionally oriented physicians find that specific botanicals are highly effective for treating infections.

BERBERIS VULGARIS has been used for centuries for lung, spleen, and liver health. *Berberis* is used for support of the urinary tract. Traditional uses include liver and gallbladder health and digestive support.

BERBERINE SULFATE is a concentrate of the main active ingredient in berberine-containing botanicals. As such, it intensifies the action of *Berberis vulgaris*.

UVA URSI (*uva ursi*) has its traditional use in treatment for urinary tract health. Recently, sensitivity cultures and clinical treatment have found a favorable use for *uva ursi* in intestinal dysbiosis treatment. *Uva ursi*'s active ingredients are found in its hydroquinone glycosides (arbutins), tannins, mono and triterpenes, and flavonoids.

GRAPEFRUIT SEED EXTRACT had no adverse effects on normal fecal organisms. Individuals consuming Grapefruit seed extract have experienced improvements in intestinal regularity.

REFERENCES:

1. Ubebaba K et al., *Jpn J Phamacol* 36 (Suppl): 352, 1984.
2. Liu CX et al., Chinese Traditional and Herbal Drugs Communications 9: 36, 1979.
3. Amin AH, Subbaiah TV, Abbasi KM. Berberine sulfate: antimicrobial activity, bioassay, and mode of action. *Can J Microbiol* 15 (9):1067-76, 1969
4. Sun D, Courtney HS, Beachey EH. Berberine sulfate blocks adherence of *Streptococcus pyogenes* to epithelial cells, fibronectin, and hexadecane. *Antimicrob Agents Chemother* 32:1370-4, 1988.
5. Mahajan VM, Sharma A, Rattan A. Antimycotic activity of berberine sulfate: An alkaloid from an Indian medicinal herb. *Sabouraudia* 20:79-81, 1982.
6. Gupte S. Use of berberine in the treatment of giardiasis. *Am J Dis Child* 129:866, 1975.



PARABIOTIC PLUS™

REFERENCES:

7. Kaneda Y, Torii M, et al. In vitro effects of berberine sulfate on the growth of *Entamoeba histolytica*, *Giardia lamblia* and *Trichomonas vaginalis*. *Ann Trop Med Parasitol* 85: 417-25, 1991.
8. Ionescu G et al. Oral citrus seed extract in atopic eczema: In vitro and in vivo studies on intestinal microflora. *J Orthomol Med* 5: 155-7, 1990.
9. Blackwell AL. Tea tree oil and anaerobic (bacterial) vaginosis. *Lett. Lancet* 337: 300, 1991.
10. Altman PM. Australian tea tree oil. *Australian J Pharmacy* 69: 276-8, 1988.
11. Sourgens H et al. *Planta Med* 45: 78, 1982.
12. Ng TB et al. Examination of coumarins, flavonoids and polysaccharopeptides for antibacterial activity. *In: General Pharmacology* 27(7): 1237-1240, 1996.
13. Bianco A et al., Partial synthesis of oleuropein. *In: JNP* 55(6): 760-66, 1992.
14. Afshaypuor S et al., Volatile constituents of *Origanum vulgare* ssp. *Virde* (syn *O. heracleoticum*) from Iran. *In: PM* 63(2): 179-180, 1997.
15. PDR for Herbal Medicines, 1st Ed. *Med Econ Co., Montvale, New Jersey, 1998.*
16. Werbach MR, Murray MT. Botanical Influences on Illness: A sourcebook of clinical research. *Third Line Press, Tarzana, California, 1994.*

JUGLANS NIGRA (BLACK WALNUT HULL) has a long traditional use as a vermifuge. It has been successfully employed in the management of parasitic dysbiosis.

THYMUS VULGARIS (Thyme) is often used to treat cough. Thyme is a bronchial antispasmodic, an expectorant, and an antimicrobial agent. Active ingredients include its volatile oils (thymol, p-cymene, carvacrol, borneol, linalool), caffeic acid, flavonoids, and triterpenes.

ORIGANUM VULGARE (Oregano) has often been used to treat respiratory disorders and complaints such as coughs and bronchial congestion.

OLEA EUROPAEA EXTRACT (Olive leaf) is reported to have treated over 650 pathogens successfully. Olive leaf extract is antiviral and antimicrobial. In animal studies it enhanced coronary circulation. Active ingredients include its mono and triterpenes, and flavonoids.

ARTEMESIA ANUUA EXTRACT (Wormwood) possesses antimicrobial effects. Artemesia is often used to treat loss of appetite, dyspepsia, liver/gallbladder complaints, intestinal bloating, gastric insufficiency, gastritis, stomach ache, and dysbiosis.

Precautions: This formula is not recommended for use during pregnancy, by nursing mothers, or by children under the age of 12.

*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.